Migraleve Pink

Uses

For the short term treatment of acute moderate pain which is not relieved by paracetamol, ibuprofen or aspirin alone, such as migraine attacks including the symptoms of migraine headache, nausea and vomiting. Not to be taken for more than 3 days continuously without medical review. Can cause addiction.

Dosage

Do not take for more than 3 days continuously without medical review. Can cause addiction. 16 years and over: 2 Pink tablets at onset. If required, 2 Yellow tablets every 4 hours. Maximum 2 Pink and 6 Yellow tablets in 24 hours. 12-15 years: 1 Pink tablet at onset. If required 1 Yellow tablet 4 hourly. Maximum 1 Pink and 3 Yellow tablets in 24 hours. Under 12 years: not to be given.

Format and Ingredients

Pink film-coated tablet containing:
Paracetamol 500mg, Codeine phosphate 8mg, Buclizine Hydrochloride 6.25mg,

Category:
Analgesics

Manufacturer:
KENVUE

Pack size: 12, 24

RRP: £9.10, £14.45

Legal Status: (P), (P)

A division of GROUPE EUROCOM
PAGB
Communications International Group

Product data © Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB